



TIME-INDEPENDENT PERTURBATIONS

Quantum Physics

Project PHYSNET Physics Bldg. Michigan State University East Lansing, MI

TIME-INDEPENDENT PERTURBATIONS

by
R. Spital

1. Introduction	1
2. Procedures	1
Acknowledgments.....	2

Title: **Time-Independent Perturbations**

Author: R. Spital, Dept. of Physics, Illinois State Univ

Version: 2/1/2000

Evaluation: Stage B0

Length: 2 hr; 8 pages

Input Skills:

1. Vocabulary: diagonal matrix elements, off-diagonal matrix elements.
2. Unknown: assume (MISN-0-389).

Output Skills (Knowledge):

- K1. Given any time-independent perturbation \mathcal{H}_1 and any complete set of non-degenerate eigenstates of \mathcal{H}_0 , derive: (a) the energy shifts to first and second order in \mathcal{H}_1 , (b) the corrections to the wave-functions to first and second order in \mathcal{H}_1 ; and using the results of (a) and (b) above, write approximate expressions for the eigenvalues and eigenfunctions of \mathcal{H} to first or second order in \mathcal{H}_1 .

Output Skills (Problem Solving):

- S1. Solve problems such as 9-5, 9-9, 9-10, 9-11 and 9-13.

External Resources (Required):

1. E. E. Anderson, *Modern Physics and Quantum Mechanics*, W. B. Saunders (1971).

THIS IS A DEVELOPMENTAL-STAGE PUBLICATION
OF PROJECT PHYSNET

The goal of our project is to assist a network of educators and scientists in transferring physics from one person to another. We support manuscript processing and distribution, along with communication and information systems. We also work with employers to identify basic scientific skills as well as physics topics that are needed in science and technology. A number of our publications are aimed at assisting users in acquiring such skills.

Our publications are designed: (i) to be updated quickly in response to field tests and new scientific developments; (ii) to be used in both classroom and professional settings; (iii) to show the prerequisite dependencies existing among the various chunks of physics knowledge and skill, as a guide both to mental organization and to use of the materials; and (iv) to be adapted quickly to specific user needs ranging from single-skill instruction to complete custom textbooks.

New authors, reviewers and field testers are welcome.

PROJECT STAFF

Andrew Schnepf	Webmaster
Eugene Kales	Graphics
Peter Signell	Project Director

ADVISORY COMMITTEE

D. Alan Bromley	Yale University
E. Leonard Jossem	The Ohio State University
A. A. Strassenburg	S. U. N. Y., Stony Brook

Views expressed in a module are those of the module author(s) and are not necessarily those of other project participants.

© 2001, Peter Signell for Project PHYSNET, Physics-Astronomy Bldg., Mich. State Univ., E. Lansing, MI 48824; (517) 355-3784. For our liberal use policies see:

<http://www.physnet.org/home/modules/license.html>.

TIME-INDEPENDENT PERTURBATIONS

by
R. Spital

1. Introduction

Relatively few Hamiltonians are solvable exactly; most of the time we have to find ways to approximate the energy eigenstates and eigenvalues. Often we can write the Hamiltonian as $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_1$, where \mathcal{H}_0 has *known* eigenfunctions and eigenvalues, while \mathcal{H}_1 (which contains the new physics) can be regarded as small compared to \mathcal{H}_0 . The procedure for approximating the eigenstates and eigenvalues of \mathcal{H} for the case when \mathcal{H}_1 is independent of time is called “Time Independent Perturbation Theory.”

2. Procedures

1. Read section 9.1. Follow the derivations carefully and be sure you can reproduce them. The key results to remember are equations (9.8), (9.10), (9.12), (9.13) and (9.14).

Two key assumptions enter this derivation:

- (a) The eigenstates of \mathcal{H}_0 are *non-degenerate*. If this were not true, some of the “energy denominators” in the above equations would vanish, leading to nonsensical results. The Perturbation theory of degenerate states is a more advanced topic which we shall not have time to take up.
 - (b) The eigenstates of \mathcal{H}_0 and of \mathcal{H} are in one to one correspondence. This is *not* always true, and when it isn’t things are much more complicated. We shall always make this assumption.
- Having finished the derivations, equation (9.5) tells you how to write the approximate expressions for the energies and the eigenstates.
2. Solve problems 9-5, 9-9, 9-10, 9-11, and 9-13 to gain practice in applying the newly developed techniques. You may wish to read section 9.2 to see an example done, but this is not required.

Acknowledgments

The author would like to thank Illinois State University for support in the construction of this lesson. Preparation of this module was supported in part by the National Science Foundation, Division of Science Education Development and Research, through Grant #SED 74-20088 to Michigan State University.

